


Department of Defense  
Armed Forces Health Surveillance Branch  
Global Zika Virus Surveillance Summary  
(28 DEC 2016)



**Approved for Public Release**

*For questions or comments, please contact:*

[dha.ncr.health-surv.list.afhs-ib-alert-response@mail.mil](mailto:dha.ncr.health-surv.list.afhs-ib-alert-response@mail.mil)



# DEPARTMENT OF DEFENSE (AFHSB)

## Global Zika Virus Surveillance Summary #50

### 28 DEC 2016



**DoD SURVEILLANCE:** Weekly incidence among Military Health System (MHS) beneficiaries has decreased significantly since its peak during the week ending 30 JUL 2016. As of 1300 on 28 DEC, there have been 161 confirmed Zika virus (ZIKV) disease cases (see table) since the first case was reported during the third week of 2016. There are four cases in pregnant Service members and one case in a pregnant dependent.

On 7 DEC, AFHSB issued [updated detection and reporting guidance](#) that includes delineation of ZIKV virus infection versus ZIKV disease case definitions with reporting information for each. Cases should be reported in DRSi as “Any Other Unusual Condition Not Listed,” with “Zika” entered in the comment field along with additional pertinent information such as travel history and pregnancy status.

IgM ELISA and rRT-PCR assays are available under an [Emergency Use Authorization \(EUA\)](#) at DoD laboratories (see map on [Slide 4](#)). Confirmatory PRNT testing is available at the NIDDL.

As of 31 OCT, no vector mosquitoes collected on DoD installations had tested positive for ZIKV.

**CASE REPORT:** Overall weekly incidence for travel-associated cases in the U.S. States and locally-acquired cases in Puerto Rico are trending downwards.

As of 27 DEC, TX has reported six (+1) cases of suspected locally transmitted ZIKV disease in Cameron County, TX, since it announced its first ZIKV case likely transmitted by a mosquito on 28 NOV. The sixth case is not associated with the first five cases. Cameron County is in southeast TX and borders the Mexican state of Tamaulipas, which has reported 69 ZIKV cases in 2016.

As of 28 DEC, FL health officials have reported 256 (+4) locally acquired ZIKV infections. On DEC 9, FL announced the end of sustained active transmission in Miami-Dade County and sporadic cases may continue to occur. FL DOH continues

Zika Cases in the U.S. States and Territories	U.S. States*	U.S. Territories		
		Puerto Rico**	U.S. Virgin Islands*	American Samoa*
Total Zika Cases	4,756 (+139)	35,870 (+222)	877 (+13)	114 (+42)
Travel-Associated***	4,541 (+110)	-	-	-
Local Vector Transmission	215 (+30)	-	-	-
Laboratory Exposure	1	-	-	-
Guillain Barré Syndrome (GBS)	13	66 (-1)†	-	-

U.S. Zika Pregnancy Registry Data, as of 13 DEC		
Pregnant Zika Cases	1,246 (+74)	2,701 (+62)
Infants Born with Birth Defects	34 (+2)	1††
Pregnancy Losses with Birth Defects	5	1††

\*Zika cases reported to ArboNET as of 21 DEC (U.S. States and Am. Samoa). Zika cases reported by USVI as of 27 DEC; USVI also reported 106 (+2) Zika cases in pregnant women.

\*\*From the Puerto Rico DOH as of 8 DEC; PR DOH is tracking 2,880 (+16) ZIKV cases in pregnant women.

\*\*\*Includes 38 sexually transmitted cases.

† Of the 66 (-1) GBS cases, 16 are classified as evidence of flavivirus infection, but specific virus undetermined.

†† CDC last reported these cases on 29 SEP.

Demographics for all confirmed Zika cases in Military Health System Beneficiaries as of 1300, 28 DEC 2016 (N = 161 confirmed cases)			
Demographic		N	%
<b>Service</b> <small>*includes MHS beneficiaries from USPHS, NOAA, etc.</small>	Army	70	43.5%
	Air Force	27	16.8%
	Navy	20	12.4%
	Marine Corps	12	7.5%
	Coast Guard	30	18.6%
	Other*	2	1.2%
<b>Status</b> <small>**includes Reserve Component</small>	Service Member**	113	70.2%
	Dependent	37	23.0%
	Retiree	11	6.8%
<b>Age</b>	0-20	11	6.8%
	21-35	76	47.2%
	36-50	48	29.8%
	51+	19	11.8%
	Not Reported	7	4.3%
<b>Gender</b>	Female	64	39.8%
	Male	97	60.2%

to investigate additional areas in Miami-Dade County.

Updated advice for people living in or traveling to Miami-Dade County, FL, and Cameron County, TX, is available from [CDC](#).

[FL DOH and CDC](#) said aggressive mosquito control, including aerial spraying that targeted adult and larval mosquitoes, most likely contributed to stopping ZIKV transmission in the Wynwood neighborhood.

CDC has issued Alert Level 2, Practice Enhanced Precautions, travel notices for 60 [countries and territories](#); 49 are in the Western Hemisphere, 10 are in PACOM, and one is in AFRICOM.

[CDC has posted travel information](#) for 11 countries in Southeast Asia. The countries are: Brunei, Burma (Myanmar), Cambodia, Indonesia, Laos, Malaysia, Maldives, Philippines (52 (+13) cases), Thailand (>680 cases), Timor-Leste (East Timor), and Vietnam (155 (+52) cases).

(+xx) represent the change in number from the previous AFHSB summary (21 DEC 2016).

All information has been verified unless noted otherwise.

For questions or comments, please contact: [dha.ncr.health-surv.list.afhs-ib-alert-response@mail.mil](mailto:dha.ncr.health-surv.list.afhs-ib-alert-response@mail.mil)

**Approved for Public Release**



# DEPARTMENT OF DEFENSE (AFHSB)

## Global Zika Virus Surveillance Summary #50

### 28 DEC 2016



**CASE REPORT (cont'd):** These countries have either reported low level local ZIKV transmission or are adjacent to countries with known ZIKV transmission. Singapore continues to report low-level ZIKV transmission with 457 cases and no identified clusters as of 21 DEC. Past evidence of local transmission has been reported from other areas of [Africa, Asia, and the Pacific Islands](#), where sporadic transmission may continue to occur. According to [CDC](#), increased case reporting from PACOM countries, some of which are endemic for ZIKV, may be the result of increased testing and surveillance or a change in the intensity of virus transmission.

According to [PAHO](#) on 15 DEC, over the previous four weeks nearly all Caribbean and North, Central, and South American OCONUS countries and territories reported a decreasing trend in Zika cases except for Panama, Anguilla, Paraguay, and the Iquitos region of Peru.

**MICROCEPHALY and GUILLAIN-BARRÉ SYNDROME:** According to [WHO](#), as of **21 DEC**, 29 countries have reported cases of microcephaly and other fetal malformations potentially associated with ZIKV infection or suggestive of a congenital infection, including four with travel-related microcephaly cases. As of **21 DEC**, 20 countries and territories in the Western Hemisphere as well as French Polynesia have reported Guillain-Barré syndrome (GBS) cases that may be associated with of ZIKV infection. The Western Hemisphere countries reporting microcephaly or GBS are listed in the table on [slide 7](#). Countries in PACOM and AFRICOM reporting microcephaly are Cape Verde, French Polynesia, the Marshall Islands, Thailand, and Vietnam.

**USG RESPONSE:** On 22 DEC, [FDA issued a warning](#) to providers making them aware of higher than expected false positive results on IgM tests, especially the Zika Detect IgM Capture ELISA, and advising them to wait for confirmatory testing before making patient management decisions. [CDC announced](#) on 22 DEC that it was awarding nearly \$184 million of the \$350 million it received in the Zika Response and Preparedness Appropriations Act of 2016 to states, territories, local jurisdictions, and universities to support the response to ZIKV infection and its associated adverse health outcomes. On 16 NOV, CDC released [Updated: Guidance for US Laboratories Testing for Zika Virus Infection](#). CDC issued [ZIKV infection control guidance](#) on 25 OCT. On 30 SEP, [CDC updated its interim guidance](#) for pre-conception counseling and for preventing sexual transmission of ZIKV among exposed persons. The primary change was a recommendation that men with possible ZIKV exposure, but no symptoms, wait six months after the last possible ZIKV exposure before attempting conception with their partner. Also on 30 SEP, CDC published an updated [ZIKV response plan for CONUS and Hawaii](#).

**GLOBAL RESPONSE:** Following the fifth meeting of the Emergency Committee (EC) on ZIKV, microcephaly, and other neurological disorders on 18 NOV, WHO declared that the [event no longer meets the criteria](#) for a Public Health Emergency of International Concern (PHEIC). The EC said that ZIKV and its associated consequences remain a significant enduring public health challenge requiring intense action, but is no longer a PHEIC as defined under the International Health Regulations. WHO had declared the PHEIC on 1 FEB 2016. On 25 OCT, WHO issued the [first quarterly update](#) to its [JUL 2016 Zika Strategic Response Plan](#). PAHO has created a [searchable database](#) of published primary research and protocols. For additional information, visit the [WHO](#) and [PAHO](#) Zika web pages.

**MEDICAL COUNTERMEASURES and RESEARCH:** A study published in PLOS One on 20 DEC describing the prevalence of ZIKV antibodies in mothers from Hawaii who gave birth to babies with and without microcephaly between 2009 and 2012 suggests the presence of ZIKV infections and associated microcephaly in the U.S. as early as 2009. On 21 DEC, Inovio Pharmaceuticals announced its DNA-based Zika vaccine (GLS-5700) generated robust antigen-specific antibody responses in its first multi-center phase I trial. [EID posted a case series](#) from Guadeloupe (seven cases) suggesting a casual link between severe thrombocytopenia and ZIKV infection. In an [early release Emerging Infectious Diseases \(EID\) article](#), researchers found that ZIKV can replicate and persist in fetal brain and placental tissue for months after maternal infection. On 14 DEC, researchers published a study in JAMA using preliminary data from the U.S. Pregnancy Registry. This study showed that 6% of fetuses or infants had ZIKV-associated birth defects among pregnant women with evidence of recent ZIKV infection who completed pregnancies; no birth defect cases were attributed to maternal exposure solely in the second or third trimester. On 13 DEC, researchers published a cohort study in NEJM characterizing the spectrum of fetal outcomes among ZIKV infected pregnant women in Brazil; adverse fetal outcomes were evident regardless of the trimester of maternal infection (55% of pregnancies had adverse outcomes after maternal infection during the first trimester, 52% after infection during the second trimester, and 29% after infection during the third trimester). In an [early release MMWR article](#) published on 9 DEC, researchers studying ZIKV disease among pregnant women in Colombia found the risk of ZIKV related birth defects was highest when maternal infection occurred during the first trimester or early second trimester of pregnancy; evidence also suggests ZIKV related birth defects in Colombia may be significantly underreported. The Walter Reed Army Institute of Research (WRAIR) began [Phase 1 clinical testing](#) of a Zika purified inactivated virus (ZPIV) on 7 NOV. On 3 NOV, JAMA Pediatrics published a review of the distinctive features of congenital Zika syndrome in infants. In an [early release EID article](#), researchers estimated the incidence of GBS in Puerto Rico following the introduction of ZIKV was 3.2 to 5.1 times above baseline in 2016. On 17 OCT, EID posted research showing that ZIKV RNA could be isolated in [vaginal secretions, whole blood, and semen](#) up to 14 days, 81 days, and 92 days after symptom onset, respectively. The authors in both reports caution that the detection of ZIKV RNA does not necessarily equate to the detection of infectious virus.

(+xx) represent the change in number from the previous AFHSB summary (21 DEC 2016).

All information has been verified unless noted otherwise.

For questions or comments, please contact: [dha.ncr.health-surv.list.afhsb-ib-alert-response@mail.mil](mailto:dha.ncr.health-surv.list.afhsb-ib-alert-response@mail.mil)




# DEPARTMENT OF DEFENSE (AFHSB)

## Global Zika Virus Surveillance Summary #50

28 DEC 2016



### Emergency Use Authorization Zika Testing at DoD Laboratories



- BAMC**  
Brooke Army Medical Center
- BAACH**  
Brian Allgood Army Community Hospital
- CRDAMC**  
Carl R. Darnall Army Medical Center
- EAMC**  
Eisenhower Army Medical Center
- LRMC**  
Landstuhl Regional Medical Center
- MAMC**  
Madigan Army Medical Center
- NAMRU-3**  
U.S. Naval Medical Research Unit No. 3
- NAMRU-6**  
U.S. Naval Medical Research Unit No. 6
- NHRC**  
Naval Health Research Center
- NIDDL**  
Naval Infectious Diseases Diagnostic Laboratory
- TAMC**  
Tripler Army Medical Center
- USAFSAM**  
U.S. Air Force School of Aerospace Medicine
- USAMRIID**  
United States Army Medical Research Institute of Infectious Diseases
- WAMC**  
Womack Army Medical Center
- WBAMC**  
William Beaumont Army Medical Center
- WRNMMC**  
Walter Reed National Military Medical Center

\*Plaque-reduction neutralization test (PRNT)

As of 28 DEC 2016

For questions or comments, please contact: [dha.ncr.health-surv.list.afhs-ib-alert-response@mail.mil](mailto:dha.ncr.health-surv.list.afhs-ib-alert-response@mail.mil)


Approved for Public Release

# DEPARTMENT OF DEFENSE (AFHSB)


## Global Zika Virus Surveillance Summary #50

### 28 DEC 2016

### Zika Virus Distribution as of 28 DEC 2016



- Countries/states with active ZIKV transmission since 2015
- Endemic countries currently reporting ZIKV transmission and adjacent countries
- Endemic countries currently not reporting an outbreak



\* Countries with a small footprint are given a marker by their label to denote current or previous Zika presence. Source: CDC.


For questions or comments, please contact: [dha.ncr.health-surv.list.afhs-ib-alert-response@mail.mil](mailto:dha.ncr.health-surv.list.afhs-ib-alert-response@mail.mil)



# DEPARTMENT OF DEFENSE (AFHSB)

## Overlap of States Reporting Imported Zika Cases with the Estimated Range of Mosquito Vectors and Transmission Suitability

28 DEC 2016



APPROVED FOR PUBLIC RELEASE

This version of the map shows that after JUL the northern extent begins to move southward.

Based on Sang et al, Predicting Unprecedented Dengue Outbreak Using Imported Cases and Climatic Factors in Guangzhou, 2014. PLoS Negl Trop Dis 9(5);e0003808.

For questions or comments, please contact: [dha.ncr.health-surv.list.afhs-ib-alert-response@mail.mil](mailto:dha.ncr.health-surv.list.afhs-ib-alert-response@mail.mil)

Approved for Public Release



# DEPARTMENT OF DEFENSE (AFHSB)

## Global Zika Virus Surveillance Summary #50

### 28 DEC 2016



#### Western Hemisphere Countries<sup>‡</sup> and Territories with Autochthonous Transmission of Zika Virus: 1 JAN 2015 – 22 DEC 2016

	Confirmed	Suspected	Microcephaly Cases*	Reporting GBS <sup>†</sup>
<b>Total</b>	176,258	533,732	2,408	20 Countries/Territories

Country/Territory	Confirmed	Suspected	Microcephaly Cases*	Reporting GBS <sup>†</sup>
Anguilla	8	58		
Antigua & Barbuda	14	465		
Argentina	26	1,821	1	
Aruba	28	676		
Bahamas	22	0		
Barbados	46	699		
Belize	66	715		
Bolivia	140	741	9	Yes
Bonaire, St. Eustatius, Saba	85	0		
Brazil	109,596	211,770	2,228	Yes
British Virgin Islands	52	74		
Cayman Islands	30	211		
Colombia	8,826	97,136	72	Yes
Costa Rica	1,581	3,840	2	Yes
Cuba	3	0		
Curaçao	820	0		
Dominica	79	1,150		
Dominican Republic	331	4,903	22	Yes
Ecuador	853	2,678		Yes
El Salvador	51	11,401	4	Yes
French Guiana	483	9,700	14	Yes
Grenada	111	314	1	Yes
Guadeloupe	379	30,845	1	Yes
Guatemala	788	3,343	15	Yes

Country/Territory	Confirmed	Suspected	Microcephaly Cases*	Reporting GBS <sup>†</sup>
Guyana	37	0		
Haiti	5	2,955	1	Yes
Honduras	298	31,936	2	Yes
Jamaica	186	7,052		Yes
Martinique	12	36,680	14	Yes
Mexico	7,335	0		Yes
Montserrat	5	2		
Nicaragua	2,053	0	2	
Panama	652	2,570	5	Yes
Paraguay	14	580	2	
Peru	244	1,098		
Puerto Rico	35,870	0	10	Yes
Saint Barthelemy	61	950		
Saint Kitts & Nevis	33	549		No
Saint Lucia	50	822		
Saint Martin	200	3,020		
Saint Vincent & the Grenadines	83	508		
Sint Maarten	62	168		
Suriname	723	2,758	2	Yes
Trinidad and Tobago	643	0	1	
Turks & Caicos	17	179		
U.S. Virgin Islands	877	130		
Venezuela	2,380	59,235		Yes

\* Number of microcephaly and/or CNS malformation cases suggestive of congenital infections or potentially associated with ZIKV infection

† Reported increase in GBS cases associated with the introduction of ZIKV and/or GBS case(s) linked to ZIKV infection

‡ Excludes the U.S.; this data can be found elsewhere in this report.

All data was obtained from PAHO, Ministries of Health, and Departments of Health unless otherwise noted.

For questions or comments, please contact: [dha.ncr.health-surv.list.afhs-ib-alert-response@mail.mil](mailto:dha.ncr.health-surv.list.afhs-ib-alert-response@mail.mil)

**Approved for Public Release**